

CLASS: XII

Time : 3.00 hrs.

MATHEMATICS

Max. Marks : 200

Section – A

Note: 1. Answer all the questions.

2. Choose the most suitable answer from the given four alternatives.

40 x 1 = 40

1. If $ae^x + be^y = c$; $pe^x + qe^y = d$ and $\Delta_1 = \begin{vmatrix} a & b \\ p & q \end{vmatrix}$; $\Delta_2 = \begin{vmatrix} c & b \\ d & q \end{vmatrix}$; $\Delta_3 = \begin{vmatrix} a & c \\ p & d \end{vmatrix}$ then the value of (x, y) is

- a) $\left(\frac{\Delta_2}{\Delta_1}, \frac{\Delta_3}{\Delta_1}\right)$ b) $\left(\log \frac{\Delta_2}{\Delta_1}, \log \frac{\Delta_3}{\Delta_1}\right)$ c) $\left(\log \frac{\Delta_1}{\Delta_3}, \log \frac{\Delta_1}{\Delta_2}\right)$ d) $\left(\log \frac{\Delta_1}{\Delta_2}, \log \frac{\Delta_1}{\Delta_3}\right)$

2. In a system of 3 linear non-homogeneous equation with three unknowns,

if $\Delta = 0$ and $\Delta_x = 0$, $\Delta_y \neq 0$, $\Delta_z = 0$ then the system has

- a) unique solution b) two solutions
c) infinitely many solution d) no solution.

3. If A and B are any two matrices such that $AB = 0$ and A is non-singular, then

- a) $B = 0$ b) B is singular c) B is non-singular d) B

4. In the homogeneous system $\rho(A) <$ the number of unknowns then the system has

- a) only trivial solution b) trivial solution and infinitely many non-trivial solutions
c) only non-trivial solutions d) no solution

5. The centre and radius of the sphere given by $x^2 + y^2 + z^2 - 6x + 8y - 10z + 1 = 0$

- a) $(-3, 4, -5)$, 49 b) $(-6, 8, -10)$, 1 c) $(3, -4, 5)$, 7 d) $(6, -8, 10)$,

6. The point of intersection of the lines $\frac{x-6}{-6} = \frac{y+4}{+4} = \frac{z-4}{-8}$ and $\frac{x+1}{2} = \frac{y+2}{4} = \frac{z+3}{-2}$ is

- a) $(0, 0, -4)$ b) $(1, 0, 0)$ c) $(0, 2, 0)$ d) $(1, 2, 0)$

7. The equation of the plane passing through the point $(2, 1, -1)$ and the line of intersection of the planes $\vec{r} \cdot (\vec{i} + 3\vec{j} - \vec{k}) = 0$ and $\vec{r} \cdot (\vec{j} + 2\vec{k}) = 0$ is,

- a) $x+4y-z=0$ b) $x+9y+11z=0$ c) $2x+y-z+5=0$ d) $2x-y+z=0$

8. $\vec{r} = s\vec{i} + t\vec{j}$ is the equation of

- a) a straight line joining the points \vec{i} and \vec{j} b) xoy plane
c) yoz plane d) zox plane

9. If $\vec{a}, \vec{b}, \vec{c}$ are three mutually perpendicular unit vectors, then $|\vec{a} + \vec{b} + \vec{c}| =$

- a) 3 b) 9 c) $3\sqrt{3}$ d) $\sqrt{3}$

10. The projection of $3\vec{i} + \vec{j} - \vec{k}$ on $4\vec{i} - \vec{j} + 2\vec{k}$ is

- a) $\frac{9}{\sqrt{21}}$ b) $\frac{-9}{\sqrt{21}}$ c) $\frac{81}{\sqrt{21}}$ d) $\frac{-81}{\sqrt{21}}$

11. The value of $i + i^{22} + i^{23} + i^{24} + i^{25}$ is

- a) i b) $-i$ c) 1 d) -1

12. If $\frac{1-i}{1+i}$ is a root of $ax^2 + bx + 1 = 0$, where a, b are real then (a,b) is

- a) $(1,1)$ b) $(1,-1)$ c) $(0,1)$ d) $(1,0)$

CENTUM ASSURED MODEL PUBLIC EXAM 2

WWW.MATHSTIMES.COM

26. The area bounded by the parabola $y^2 = x$ and its latus rectum is.
 a) $4/3$ b) $1/6$ c) $2/3$ d) $8/3$
27. The area of the region bounded by the graph of $y = \sin x$ and $y = \cos x$ between $x = 0$ and $x = \pi/4$ is.
 a) $\sqrt{2} + 1$ b) $\sqrt{2} - 1$ c) $2\sqrt{2} + 1$ d) $2\sqrt{2} + 2$
28. $\int_a^b f(x) dx =$
 a) $2 \int_0^a f(x) dx$ b) $\int_a^b f(a-x) dx$ c) $\int_a^b f(b-x) dx$ d) $\int_a^b f(a+b-x) dx$
29. Integrating factor of $\frac{dy}{dx} + \frac{1}{x \log x} \cdot y = \frac{2}{x^2}$ is
 a) e^x . b) $\log x$. c) $1/x$ d) e^{-x} .
30. The differential equation of the family of lines $y = mx$ is.
 a) $\frac{dy}{dx} = m$ b) $y dx - x dy = 0$ c) $\frac{d^2y}{dx^2} = 0$ d) $y dx + x dy = 0$
31. The complementary function of $(D^2 + 1) y = e^{2x}$ is.
 a) $(Ax+B)e^x$. b) $A \cos x + B \sin x$. c) $(Ax+B)e^{2x}$. d) $(Ax+B)e^{-x}$.
32. The order and degree of the differential equation are $\frac{d^2y}{dx^2} - y + \left(\frac{dy}{dx} + \frac{d^3y}{dx^3}\right)^{\frac{3}{2}} = 0$
 a) 2,3 b) 3,3 c) 3,2 d) 2,2
33. Which of the following is a tautology ?
 a) $p \vee q$. b) $p \wedge q$ c) $p \vee \sim p$. d) $p \wedge \sim p$.
34. Which of the following is not a binary operation on R ?
 a) $a * b = ab$. b) $a * b = a - b$ c) $a * b = \sqrt{ab}$ d) $a * b = \sqrt{a^2 + b^2}$
35. The order of $-i$ in the multiplicative group of 4th roots of unity
 a) 4 b) 3 c) 2 d) 1
36. If p is true and q is false then which of the following statements is not true ?
 a) $p \rightarrow q$ is false b) $p \vee q$ is true c) $p \wedge q$ is false d) $p \leftrightarrow q$ is true
37. A random variable X has the following p.d.f.
- | | | | | | | | | |
|----------|---|-----|------|------|------|-------|--------|----------|
| x | 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| $P(X=x)$ | 0 | k | $2k$ | $2k$ | $3k$ | k^2 | $2k^2$ | $7k^2+k$ |
- The value of k is
 a) $1/8$ b) $1/10$ c) 0 d) -1 or $1/10$
38. The mean of binomial distribution is 5 and its standard deviation is 2. Then the value of n and p are.
 a) $\left(\frac{4}{5}, 25\right)$ b) $\left(25, \frac{4}{5}\right)$ c) $\left(\frac{1}{5}, 25\right)$ d) $\left(25, \frac{1}{5}\right)$
39. If a random variable X follows Poisson distribution such that $E(X^2) = 30$ then the variance of the distribution is.

CENTUM ASSURED MODEL PUBLIC EXAM 2

WWW.MATHSTIMES.COM

- a) 6 b) 5 c) 30 d) 25

40. The p.d.f of the standard normal variate Z is $\phi(z)=$

- a) $\frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}z^2}$ b) $\frac{1}{\sqrt{2\pi}} e^{-z^2}$ c) $\frac{1}{\sqrt{2\pi}} e^{\frac{1}{2}z^2}$ d) $\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$

Section – B

Note: 1. Answer any 10 questions.

10×6=60

2. Question No. 55 is compulsory and choose any nine questions from the remaining

41. Solve the following non-homogeneous equation of three unknowns.

$$2x + 2y + z = 5 ; x - y + z = 1 ; 3x + y + 2z = 4$$

42. If $A = \begin{bmatrix} 5 & 2 \\ 7 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -1 \\ -1 & 1 \end{bmatrix}$ verify that $(AB)^{-1} = B^{-1}A^{-1}$

43. Show that the following two lines are skew lines:

$$\vec{r} = (3\vec{i} + 5\vec{j} + 7\vec{k}) + t(\vec{i} - 2\vec{j} + \vec{k}) \text{ and } \vec{r} = (\vec{i} + \vec{j} + \vec{k}) + s(7\vec{i} + 6\vec{j} + 7\vec{k})$$

44. P represents the variable complex number z. Find the locus of P, if $|z - 3i| = |z + 3i|$

45. Solve: $x^4 + 4 = 0$

46. Find the angle between the asymptotes to the hyperbola $3x^2 - 5xy - 2y^2 + 17x + y + 14 = 0$

47. Prove that the curve $2x^2 + 4y^2 = 1$ and $6x^2 - 12y^2 = 1$ cut each other at right angles.

48. Evaluate: $\lim_{x \rightarrow 0} \left(\operatorname{cosec} x - \frac{1}{x} \right)$

49. Find $\frac{\partial w}{\partial u}$ and $\frac{\partial w}{\partial v}$ if $w = x^2 + y^2$ where $x = u^2 - v^2$, $y = 2uv$

50. Evaluate: $\int_{\pi/6}^{\pi/3} \frac{dx}{1 + \sqrt{\tan x}}$

51. Construct the truth tables for the following statements;

- (i) $\sim(p \vee q)$ (ii) $\sim(p \vee (\sim q))$

52. Construct truth table for $(p \vee q) \wedge r$

53. Suppose that the probability of suffering a side effect from a certain vaccine is 0.005. If 1000 persons are inoculated. Find approximately the probability that

- (i) atmost 1 person suffer. (ii) 4,5 or 6 persons suffer. $[e^{-5} = 0.0067]$.

54 For the p.d.f $f(x) = \begin{cases} cx(1-x)^3, & 0 < x < 1 \\ 0, & \text{elsewhere} \end{cases}$

- find (i) the constant C (ii) $P\left(x < \frac{1}{2}\right)$

55. (a) Prove that $\left[\vec{a} \times \vec{b}, \vec{b} \times \vec{c}, \vec{c} \times \vec{a} \right] = \left[\vec{a}, \vec{b}, \vec{c} \right]^2$ (OR)

CENTUM ASSURED MODEL PUBLIC EXAM 2

WWW.MATHSTIMES.COM

(b) Solve $(D^2 - 6D + 9)y = e^{3x}$

Section - c**Note: 1. Answer any 10 questions.****10×10=100****2. Question No. 70 is compulsory and choose any nine questions from the remaining**

56. A bag contains 3 types of coins namely Re.1, Rs. 2 and Rs. 5. There are 30 coins amounting to Rs. 100 in total. Find the number of coins in each category.

57. Prove that $\sin(A - B) = \sin A \cos B - \cos A \sin B$.58. Find the vector and Cartesian equations of the plane passing through the points (-1,1,1) and (1,-1,1) and perpendicular to the plane $x + 2y + 2z = 5$ 59. Solve the equation $x^7 + x^4 + x^3 + 1 = 0$ 60. Find the axis, vertex, focus, equation of directrix, latus rectum, length of the latus rectum for the following parabola and hence sketch their graph. $x^2 - 6x - 12y - 3 = 0$ 61. A comet is moving in a parabolic orbit around the sun which is at the focus of a parabola. When the comet is 80 million kms from the sun, the line segment from the sun to the comet makes an angle of $\frac{\pi}{3}$ radians with the axis of the orbit. Find (i) the equation of the comet's orbit (ii) how close does the comet nearer to the sun? (Take the orbit as open rightward).

62. Find the dimensions of the rectangle of largest area that can be inscribed in a circle of radius r.

63. If the curve $y^2 = x$ and $xy = k$ are orthogonal then prove that $64k^4 = 1$ 64. Verify Euler's theorem for $f(x, y) = \frac{1}{\sqrt{x^2 + y^2}}$ 65. Find the area of the region bounded by the parabola $y^2 = 4x$ and the line $2x - y = 4$.66. The sum of Rs. 1000 is compounded continuously, the nominal rate of interest being four percent per annum. In how many years will the amount be twice the original principal? ($\log_e 2 = 0.6931$)67. solve: $dx + xdy = e^{-y} \sec^2 y dy$ 68. Show that the set M of complex numbers z with the condition $|z| = 1$ forms a group with respect to the operation of multiplication of complex numbers.69. Find c , μ and σ^2 of the normal distribution whose probability function is given by

$$f(x) = ce^{-x^2 + 3x} \quad -\infty < X < \infty.$$

70. (a) Find the equation of the hyperbola if its asymptotes are parallel to $x + 2y - 12 = 0$ and $x - 2y + 8 = 0$, (2,4) is the centre of the hyperbola and it passes through (2,0).**(OR)**(b) Show that the surface area of the solid obtained by revolving the arc of the curve $y = \sin x$ from $x = 0$ to $x = \pi$ about x-axis is $2\pi \left[\sqrt{2} + \log(1 + \sqrt{2}) \right]$

WWW.MATHSTIMES.COM